AURA: Alfred University Research & Archives

https://aura.alfred.edu/

College of Liberal Arts and Sciences

Faculty Scholarship

1986

Test2 Title

Test2, Test2

Test Description Downloaded from AURA: Alfred University Research & Archives



DOI: 10.1002/jgt.22763

ARTICLE

WILEY

Bounding the tripartite-circle crossing number of complete tripartite graphs

Charles Camacho¹ | Silvia Fernández-Merchant² | Marija Jelić Milutinović³ | Rachel Kirsch⁴ | Linda Kleist⁵ Elizabeth B. Matson⁶ | Jennifer White⁷

¹University of Washington, Seattle, Washington, USA

²Department of Mathematics, California State University, Northridge, California, USA

³Faculty of Mathematics, University of Belgrade, Belgrade, Serbia

⁴Department of Mathematical Sciences, George Mason University, Fairfax, Virginia, USA

⁵Department of Computer Science, Technische Universität Braunschweig, Braunschweig, Germany

⁶Division of Mathematics and Computer Science, Alfred University, Alfred, New York, USA

⁷Department of Mathematics, Saint Vincent College, Latrobe, Pennsylvania, USA

Correspondence

Linda Kleist, Department of Computer Science, Technische Universität Braunschweig, Braunschweig 38106, Germany. Email: kleist@ibr.cs.tu-bs.de

Funding information

National Science Foundation, Grant/ AwardNumbers: 1641020, 1400653, 1839918; Ministry of Education, Science and Technological Development of Serbia, Grant/Award Number: 174034

Abstract

A tripartite-circle drawing of a tripartite graph is a drawing in the plane, where each part of a vertex partition is placed on one of three disjoint circles, and the edges do not cross the circles. We present upper and lower bounds on the minimum number of crossings in tripartite-circle drawings of $K_{m,n,p}$ and the exact value for $K_{2,2,n}$. In contrast to 1- and 2-circle drawings, which may attain the Harary–Hill bound, our results imply that balanced restricted 3-circle drawings of the complete graph are not optimal.

K E Y W O R D S

3-circle drawing, circle drawing, complete tripartite graph, crossing number, Harary–Hill bound

© 2021 The Authors. Journal of Graph Theory published by Wiley Periodicals LLC

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

1 | INTRODUCTION

The crossing number of a graph G, denoted by cr(G), is the minimum number of edge crossings over all drawings of G in the plane. It quantifies how close or far a graph is from being planar. Drawings with few crossings have been studied in connection with read-ability and VLSI chip design [23]. See [28] for a survey of crossing number variants and some of their applications. Computing the crossing number of a graph is an NP-hard problem [15,21]. The precise values are not known even for very special graph classes, such as complete and complete bipartite graphs. Nevertheless, there exist long-standing conjectures. Zarankiewicz [29] conjectured that for the complete bipartite graph $K_{m,n}$, the bound

$$\operatorname{cr}(K_{m,n}) \leq \left\lfloor \frac{m}{2} \right\rfloor \left\lfloor \frac{m-1}{2} \right\rfloor \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor =: Z(m, n),$$

given by a certain straight-line drawing of $K_{m,n}$ with vertices placed along two axes, is the best possible. Later, Harary and Hill [20] conjectured that the upper bound for the complete graph K_n ,

$$\operatorname{cr}(K_n) \leq \frac{1}{4} \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor \left\lfloor \frac{n-2}{2} \right\rfloor \left\lfloor \frac{n-3}{2} \right\rfloor =: H(n),$$

given by Guy [18], is the best possible.

Among the best-known drawings of complete graphs are drawings where the vertices are placed on one or two circles and edges do not cross the circles. Such drawings are 1-circle drawings (or 2-page book drawings) [3] and 2-circle drawings (or cylindrical drawings) [5], respectively. (For more details, refer to Section 4.) A question of interest [6,22] is to determine which other families of drawings of K_n achieve the conjectured minimum number of crossings, H(n). One possible direction is to look at greater numbers of circles.

As a natural extension of 1- and 2-circle drawings, a *k*-circle drawing of a graph *G* in the plane is a drawing in which the vertices are placed on *k* disjoint circles and the edges do not cross the circles [14]. The minimum number of crossings in a *k*-circle drawing of a graph *G* is the *k*-circle crossing number of *G*. For the special case when *G* is a *k*-partite graph, if we further require that the vertices on each circle form an independent set, we call these drawings *k*-partite-circle drawings. We call the minimum number of crossings in a *k*-partite-circle drawings he *k*-partite-circle crossing number and denote it by $cr_{\odot}(G)$. In this paper, we determine bounds for the tripartite-circle crossing number of complete tripartite graphs, and we conclude that for $n \ge 13$ there are no balanced restricted 3-circle drawings of K_n that achieve the minimum number of crossings.

1.1 | Previous results and related work

In this section, we concentrate on bipartite-circle crossing numbers and on the crossing numbers of complete tripartite graphs. The 2-circle drawings are also called cylindrical because they can be thought of as drawings on the surface of a cylinder, with the vertices on

the top and bottom circles. Analogously, a 3-circle drawing can also be understood as a pair of pants drawing (an instance of the *map crossing number* [26]), where two circles are enclosed by the third. A *radial drawing* [9] of two concentric circles is equivalent to a 2-circle drawing (cylindrical drawing). For k-circle drawings with $k \ge 3$, three or more concentric circles (or more generally three pairwise nested circles) would require that any edges from the outermost to the innermost circle would necessarily cross the middle circle (s), so a radial drawing with three or more concentric circles is not equivalent to a k-circle drawing. Consequently, 3-partite-circle drawings of complete tripartite graphs do not contain three pairwise nested circles.

The bipartite-circle drawings of bipartite graphs, in which the vertices of each part are placed on a circle and no edge crosses a circle, are of special interest due to their connection to one of the conjectured optimal drawings of K_n . A 2-circle drawing of K_n with H(n) crossings can be obtained from a bipartite-circle drawing of $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ by adding straight-line edges between vertices on the same circle. In general, the bipartite-circle crossing number of complete bipartite graphs, also known as the *bipartite cylindrical crossing number*, is fully understood. In 1997, Richter and Thomassen [27] settled the balanced case by showing that

$$\operatorname{cr}_{\odot}(K_{n,n}) = n \binom{n}{3}.$$
(1)

-WILEY

3

Ábrego, Fernández-Merchant, and Sparks [7] generalized this result to all complete bipartite graphs. For $m \le n$, the bipartite-circle crossing number is

$$\operatorname{cr}_{\odot}(K_{m,n}) = \binom{n}{2}\binom{m}{2} + \sum_{1 \le i < j \le m} \left(\left\lfloor \frac{n}{m}(j-1) \right\rfloor - \left\lfloor \frac{n}{m}(i-1) \right\rfloor \right)^{2} - n \cdot \sum_{1 \le i < j \le m} \left(\left\lfloor \frac{n}{m}(j-1) \right\rfloor - \left\lfloor \frac{n}{m}(i-1) \right\rfloor \right).$$

$$(2)$$

In particular, if *m* divides *n*, then $\operatorname{cr}_{\odot}(K_{m,n}) = \frac{1}{12} \cdot n(m-1)(2mn-3m-n)$.

For the general crossing number of complete tripartite graphs, Gethner et al. [16] proved an upper bound A(m, n, p) on $cr(K_{m,n,p})$ that is analogous to the Zarankiewicz Conjecture for $K_{m,n}$. Additionally, they proved that among *straight-line drawings* their bound is asymptotically very close to best possible. For balanced tripartite graphs, A(n, n, n) is of order ${}^{9}/_{16} \cdot n^{4}$, much less than our lower bound of ${}^{5}/_{4} \cdot n^{4}$ (see Corollary 2) because their drawings are not restricted by circles. (A similar gap exists between $Z(n, n) \sim {}^{1}/_{16} \cdot n^{4}$ and $cr_{\odot}(K_{n,n}) \sim {}^{1}/_{6} \cdot n^{4}$.) Asano [8] determined the crossing numbers of $K_{1,3,n}$ and $K_{2,3,n}$, and Ginn and Miller [17] gave bounds on $cr(K_{3,3,n})$. More recently, building upon this study, we [13] established the exact tripartite-circle crossing number of $K_{2,2,n}$ for every integer $n \geq 3$, as

$$\operatorname{cr}_{\odot}(K_{2,2,n}) = 6\left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor + 2n - 3.$$
 (3)

For other crossing number results and equivalent terminology, see, for example, [28].



1.2 | Our results

We prove several bounds on the tripartite-circle crossing number of complete tripartite graphs.

Theorem 1. Let m, n, and p be natural numbers and $\mathfrak{t} := \{(m, n, p), (n, p, m), (p, m, n)\}$. Then the following bounds hold:

$$\sum_{(a,b,c)\in\mathfrak{t}} \left(\operatorname{cr}_{\odot} (K_{a,b}) + ab \left\lfloor \frac{c-1}{2} \right\rfloor \left\lfloor \frac{c}{2} \right\rfloor \right) \leq \operatorname{cr}_{\odot}(K_{m,n,p})$$
$$\leq \sum_{(a,b,c)\in\mathfrak{t}} \left(\binom{a}{2} \binom{b}{2} + ab \left\lfloor \frac{c-1}{2} \right\rfloor \left\lfloor \frac{c}{2} \right\rfloor \right).$$

For $m, n, p \ge 3$ we improve the lower bound by 2 in Corollary 12. Using Equation (1) and Corollary 12, Theorem 1 simplifies as follows for the balanced case. Note that the lower bound of order $\sim^{5}/_{4} \cdot n^{4}$ and the upper bound of order $\sim^{6}/_{4} \cdot n^{4}$ are fairly close.

Corollary 2. For any integer $n \ge 3$,

$$3n\binom{n}{3} + 3n^{2}\left\lfloor\frac{n}{2}\right\rfloor \left\lfloor\frac{n-1}{2}\right\rfloor + 2 \le \operatorname{cr}_{\Im}(K_{n,n,n}) \le 3\binom{n}{2}^{2} + 3n^{2}\left\lfloor\frac{n}{2}\right\rfloor \left\lfloor\frac{n-1}{2}\right\rfloor$$

Finally, *k*-partite-circle drawings of complete *k*-partite graphs easily give rise to drawings of complete graphs, known as *restricted k-circle drawings*, by adding straight-line segments between each pair of vertices on the same circle. If the numbers of vertices on the circles are as close to equal as possible, these drawings are called *balanced restricted k-circle drawings* of K_n . The minimum number of crossings in such a drawing is denoted by $bcr_{\odot}(K_n)$. Certain balanced 1- and 2-circle drawings of the complete graph have H(n) crossings and are conjectured to be optimal [11,12,3,20,5]. Our results imply that this phenomenon does not generalize to balanced restricted 3-circle drawings.

Corollary 3. For n = 9, 10 and $n \ge 13$, the number of crossings in any balanced restricted 3-circle drawing of K_n exceeds H(n), that is, $bcr_{\odot}(K_n) > H(n)$.

For $n \le 7$, balanced restricted 3-circle drawings of K_n achieve the Harary–Hill bound, and we give drawings for K_6 and K_7 . For $8 \le n \le 11$, we conclude that there exist unbalanced restricted 3-circle drawings of K_n with H(n) crossings.

1.3 | Organization

The remainder of our paper is organized as follows: In Section 2, we introduce tools to count the number of the crossings, which we then use in Section 3 to prove Theorem 1 and Corollary 2. We discuss the connection to the Harary–Hill conjecture in Section 4 and conclude with a list of open problems in Section 5.

2 | TOOLS FOR COUNTING THE NUMBER OF CROSSINGS

A simple drawing of *G* is a drawing where no edge crosses itself, two edges that share a vertex do not cross, and two edges with no shared vertices intersect at most once. Drawings that minimize the number of crossings are simple, so we only consider simple drawings. In a tripartite-circle drawing of $K_{m,n,p}$, we label the three circles M, N, and P, and their numbers of vertices are *m*, *n*, and *p*, respectively. Consider Figure 1. Note that this drawing can be transformed by a projective transformation of the plane such that any one circle encloses the other two. Therefore, without loss of generality we consider drawings where the *outer* circle P contains the *inner* circles M and N. In such a drawing, we label the vertices on circles M and N in clockwise order and the vertices on circle P in counterclockwise order. Likewise, we read arcs of circles in clockwise order for inner circles and in counterclockwise order for outer circles.

2.1 | Defining the x-labels

For simplicity and without loss of generality, both papers [7,27] considered simple bipartitecircle drawings of the complete bipartite graph where the two circles are assumed to be nested. Their results rely on the assignment of a vertex x_i (A,B) on the outer circle B for each vertex *i* on the inner circle A. Because we are dealing with three circles and a pair of them is not necessarily nested, we adapt this definition as follows.

Let *i* be a vertex on circle A. The star formed by all edges from *i* to B together with circle B partitions the plane into several disjoint regions, as shown in Figure 2. Exactly one of these regions contains circle A. Such a region is enclosed by two edges from *i* to B and an arc on B between two consecutive vertices. We define the second of these vertices (in clockwise or counterclockwise order depending on whether B is an inner or outer circle, respectively) as $x_i(A,B)$. If the two circles are clear from the context, we may also write x_i .

Ábrego et al. [7] observed that the x-labels are weakly ordered and suffice to describe the drawing up to isomorphism. Because we number the vertices on the outer circle in counterclockwise order (opposite to how it is done in [7]), the ordering of x-labels on the circle is reversed when compared to Lemma 1.4 from [7]. In particular, our weak ordering stated below is achieved, following the proof from [7], by possibly renumbering the inner vertices.



FIGURE 1 The vertices on the circles M and N are labeled clockwise; the vertices of the circle P are labeled counterclockwise

5

WILFY

Lemma 4 (Ábrego et al. [7, Lemma 1.4]). Consider a simple bipartite-circle drawing of $K_{a,b}$ where the circles *A* and *B* have *a* and *b* vertices, respectively, and the vertices are labeled so that $x_1 = b$. Then it holds that

$$x_1 \ge x_2 \ge \cdots \ge x_a.$$

Moreover, for a given sequence $(s_i)_i$ with $s_1 \ge s_2 \ge \cdots \ge s_a$, up to isomorphism, there is a unique simple bipartite-circle drawing of $K_{a,b}$ with $x_i = s_i$ for all $i \in A$.

2.2 | Defining the y-labels

As observed in Lemma 4, the *x*-labels are sufficient to describe a bipartite-circle drawing. We now aim to describe tripartite-circle drawings. We therefore introduce a new vertex assignment, $y_i(A, B)$ that depends on all three circles. See Figure 3. Let A, B, and C be the three circles and *i* be a vertex on A. The star formed by all edges from *i* to B together with circle B partitions the plane into disjoint regions. Exactly one of these regions contains the third circle c. This region is enclosed by two edges incident to *i* and the arc between two consecutive vertices on B. We define the second of these two vertices (in clockwise or counterclockwise order depending on whether B is an inner or outer circle, respectively) as $y_i(A, B)$.



FIGURE 2 Definition of vertex x_i (A,B) [Color figure can be viewed at wileyonlinelibrary.com]



Nested: A is inside B

Non-Nested: A is beside B

FIGURE 3 Illustration of $y_i(A, B)$ for the cases when (a) A is inside B and (b) when A is besides B [Color figure can be viewed at wileyonlinelibrary.com]

6

2.3 Counting crossings using x- and y-labels

If two edges ab and cd cross, then at least two (nonadjacent) vertices in $\{a, b, c, d\}$ are on the same circle. Hence, there are six total types of crossings between edges ab and cd:

MP/MP-crossings:	a and c lie on M,	and b and d lie on P;
NP/NP-crossings:	a and c lie on N,	and b and d lie on P;
MN/MN-crossings:	a and c lie on M,	and b and d lie on N;
MN/MP-crossings:	a and c lie on M,	b lies on N, and d lies on P;
MN/NP-crossings:	a and c lie on N,	b lies on M, and d lies on P;
MP/NP-crossings:	a and c lie on P,	b lies on M, and d lies on N.

We typically color the edges between each pair of circles with the same color, using three different colors for the different pairs. The first three types of crossings above only involve two circles and these are called *monochromatic* crossings. The last three types involve all three circles with edges of different colors. Thus, these crossings are called *bichromatic* crossings. We use the x- and y-labels to count the monochromatic and bichromatic crossings, respectively. The following definitions are used throughout the rest of the paper.

For vertices k and ℓ on a circle with n vertices numbered 1, ..., n clockwise (respectively, counterclockwise), let

 $d_n(k, \ell) \coloneqq \ell - k \mod n$

denote the distance from k to ℓ in clockwise (respectively, counterclockwise) order on the circle. Let $[n] := \{1, 2, ..., n\}$. For any $u, v \in [n]$, define

$$f_n(u,v) \coloneqq \binom{d_n(u,v)}{2} + \binom{n-d_n(u,v)}{2}.$$

For vertices *i* and *j* on the inner (respectively, outer) circle A, we use [i, j] to denote the arc of A read clockwise (respectively, counterclockwise) from *i* to *j*. We include *i* and *j* in the interval [i, j], whereas (i, j) does not include *i* and *j*. We similarly define [i, j], and (i, j].

2.3.1 | Counting crossings involving two circles

We start by stating the following result from [27] to take care of the monochromatic crossings.

Lemma 5 (Richter and Thomassen [27, Sect. 2]). The number of crossings in a simple bipartite-circle drawing of the complete bipartite graph $K_{m,n}$ is

$$\sum_{1 \le i < j \le m} f_n(x_i, x_j).$$

2.3.2 | Counting crossings involving three circles

The following lemma introduces a means of counting all three types of bichromatic crossings using the *y*-labels. See Figure 4 for a visual representation of a possible MP/NP-crossing.

³ ⊥Wiley



FIGURE 4 Illustration for the case that (A,B,C) = (M,N,P). Since $k' \in I_1$ and $\ell \in I_2$, edges between the vertices i, j, k', and ℓ do not cross, but edges between the vertices i, j, k, and ℓ do cross since k and ℓ are in the same interval I_2 [Color figure can be viewed at wileyonlinelibrary.com]

Lemma 6. Let *A*, *B*, and *c* be three disjoint circles with the disjoint vertex sets $\{1, ..., a\}$, $\{1, ..., b\}$, and $\{1, ..., c\}$, respectively. Then the number of AC/BC-crossings is given by

$$\sum_{\substack{1 \le i \le a \\ 1 \le j \le b}} f_c(y_i(\mathbf{A}, \mathbf{C}), y_j(\mathbf{B}, \mathbf{C})).$$

Proof. Fix a vertex *i* on A and a vertex *j* on B and consider the corresponding vertices $y_i(A, C)$ and $y_j(B, C)$ on circle c, see Figure 4. For every pair of distinct vertices *k* and ℓ both in the interval $[y_i(A, C), y_j(B, C)) =: I_1$ on c there is exactly one crossing among edges *ik* and *i* ℓ , and *jk* and *j* ℓ . Similarly, there is exactly one crossing among the edges *ik* and *i* ℓ , and edges *jk* and *j* ℓ when *k* and ℓ are in $[y_j(B, C), y_i(A, C)] =: I_2$. Moreover note that if a vertex *k* is in I_1 and a vertex ℓ is in I_2 then there are no crossings among edges *ik*, *i* ℓ , *jk*, and *j* ℓ . Consequently, there are exactly $f_c(y_i(A, C), y_j(B, C))$ crossings among edges incident with vertices *i* and *j*. Therefore the total number of AC/BC-crossings is as claimed.

2.3.3 | Total crossing count

The number of crossings in a simple tripartite-circle drawing of $K_{m,n,p}$ can be found by counting the crossings in the three different, simple bipartite-circle drawings of $K_{m,n}$, $K_{m,p}$, and $K_{n,p}$, along with crossings involving all three circles. Therefore, we say a *cyclic assignment* of (A, B, C) to (M, N, P) is one triple in the set $t := \{(M,N,P), (N,P,M), (P,M,N)\}$, with the number of vertices on the circles A, B, and C denoted by *a*, *b*, and *c*, respectively.

Theorem 7. The number of crossings in a simple tripartite-circle drawing of $K_{m,n,p}$ is given by

$$\sum_{\substack{(\mathbf{A},\mathbf{B},\mathbf{C})\in\mathfrak{t}\\i,j\in\mathbf{A}}} \left(\sum_{\substack{i< j\\i,j\in\mathbf{A}}} f_b(x_i(\mathbf{A},\mathbf{B}), x_j(\mathbf{A},\mathbf{B})) + \sum_{\substack{i\in\mathbf{A}\\j\in\mathbf{B}}} f_c(y_i(\mathbf{A},\mathbf{C}), y_j(\mathbf{B},\mathbf{C})) \right)$$

Proof. The monochromatic crossings are counted by the first expressions in the brackets using Lemma 5. The second expression corresponds to the bichromatic crossings using Lemma 6.

3 | BOUNDING THE TRIPARTITE-CIRCLE CROSSING NUMBER—PROOFS OF THEOREM 1 AND COROLLARY 2

In this section, we prove the upper and lower bounds of Theorem 1 and Corollary 2. We start with the lower bounds and then proceed with the upper bounds.

3.1 | Lower bounds

To prove the lower bounds, we start with two lemmas.

Lemma 8. The function $f_n(a, b)$ attains its minimum M if and only if $|a - b| \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$. Among pairs (a, b) such that $|a - b| \notin \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$, the minimum of f_n exceeds M by 1 if n is even and by 2 if n is odd.

Proof. First note that $d_n(a, b) = \lfloor n/2 \rfloor$ if and only if $|a - b| \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$ because $n = \lfloor n/2 \rfloor + \lceil n/2 \rceil$.

Consider the auxiliary real-valued function $g_n(x) = \binom{x}{2} + \binom{n-x}{2} = \left(x - \frac{n}{2}\right)^2 + \frac{n^2 - 2n}{4}$. It is a quadratic function minimized at $\frac{n}{2}$, symmetric about $x = \frac{n}{2}$, decreasing for $x < \frac{n}{2}$, and increasing for $x > \frac{n}{2}$. Consequently, the minimum value of g_n over the integers is attained at $x \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$; the next smallest value of g_n over the integers is attained at $x \in \{\lfloor n/2 \rfloor - 1, \lceil n/2 \rceil + 1\}$. Finally, the claim follows by a computation showing that

$$g_n(\lfloor n/2 \rfloor - 1) - g_n(\lfloor n/2 \rfloor) = \begin{cases} 2 & \text{if } n \text{ is odd,} \\ 1 & \text{if } n \text{ is even.} \end{cases} \square$$

For every cyclic assignment of (A,B,C) to (M,N,P), we denote the minimum number of AB/ Ac-crossings among all simple tripartite-circle drawings of $K_{m,n,p}$ by $cr_{\odot}^{A}(K_{m,n,p})$.

Lemma 9. For every cyclic assignment of (A,B,C) to (M,N,P),

$$\operatorname{cr}^{\mathbb{C}}_{\mathfrak{S}}(K_{m,n,p}) \geq ab \left\lfloor \frac{c}{2} \right\rfloor \left\lfloor \frac{c-1}{2} \right\rfloor.$$

Proof. From the count in Lemma 6, and its minimization in Lemma 8, we have the lower bound

$$\sum_{\substack{1 \le i \le a \\ 1 \le j \le b}} f_c(y_i(\mathbf{A}, \mathbf{C}), y_j(\mathbf{B}, \mathbf{C})) \ge \sum_{\substack{1 \le i \le a \\ 1 \le j \le b}} \left| \binom{\lfloor c/2 \rfloor}{2} + \binom{\lceil c/2 \rceil}{2} = ab \left| \binom{\lfloor c/2 \rfloor}{2} + \binom{\lceil c/2 \rceil}{2} \right|.$$

Adding the last two terms directly yields the claim.

9

This allows us to find a lower bound on $\operatorname{cr}_{\odot}(K_{m,n,p})$.

Proof of the lower bound of Theorem 1. In Theorem 7 we count the total number of crossings in a simple tripartite-circle drawing of $K_{m,n,p}$.

$$\operatorname{cr}_{\odot}(K_{m,n,p}) \ge \operatorname{cr}_{\odot}(K_{m,n}) + \operatorname{cr}_{\odot}(K_{m,p}) + \operatorname{cr}_{\odot}(K_{n,p}) + \operatorname{cr}_{\odot}^{\mathrm{M}}(K_{m,n,p}) + \operatorname{cr}_{\odot}^{\mathrm{N}}(K_{m,n,p}) + \operatorname{cr}_{\odot}^{\mathrm{N}}(K_{m,n,p}).$$

Lemma 9 gives a lower bound on the last three summands and directly implies the claimed lower bound:

$$\operatorname{cr}_{\mathfrak{I}}(K_{m,n,p}) \geq \sum_{(A,B,C)\in\mathfrak{t}} \left(\operatorname{cr}_{\mathfrak{I}}(K_{a,b}) + ab \left\lfloor \frac{c}{2} \right\rfloor \left\lfloor \frac{c-1}{2} \right\rfloor \right).$$

3.1.1 | Improving the lower bound

With the help of the following lemmas, the lower bound can be improved by 2. This improvement is relevant in the analysis of small graphs and the connection to the Harary–Hill conjecture. It is also used in [13] to settle the tripartite-circle crossing number of $K_{2,2,n}$.

Lemma 10 (Special Inversion Lemma). Fix the placement of two circles A and B inside circle c. For a tripartite-circle drawing $D, x_i(A, C) = y_i(A, C)$ for all i on A if and only if $y_i(C, A) = y_k(C, A)$ for all j, k on c.

Proof. Let D^* be the restriction of D to edges between A and c. We refer to the connected components of the complement of D^* as *faces*. We start with an observation. By the definitions of $x_i(A, C)$ and $y_i(A, C)$, we have $x_i(A, C) = y_i(A, C)$ if and only if the pair of incident edges $\{i, x_i(A, C)\}$ and $\{i, x_i(A, C) + 1\}$ divides the interior of c into two parts where A and B lie in the same part and all other edges from *i* lie in the other part. This holds for all *i* if and only if no edge of D^* separates A from B; in other words, the circle B lies in a face F of D^* adjacent to A. Figure 5 shows an example with $x_i(A, C) = y_i(A, C)$ for all *i* and with the faces adjacent to A shaded in gray. Circle B lies in one of these faces, say F. This finishes our observation.

Now we prove the lemma. Suppose $x_i(A, C) = y_i(A, C)$ for all *i* on A. By the above observation, the circle B lies in a face *F* of *D*^{*} adjacent to A. Let *q* and *q* + 1 be the two vertices on A and on the boundary of *F*, where *q* + 1 comes clockwise directly after *q*. For each *j* on c, the triangle $T_j = (j, q, q + 1)$ cannot cross *F*, since it is a face, and shares the side $\{q, q + 1\}$ with *F*. By the properties of simple drawings, either *F* is in the interior of T_j and all other edges from *j* are on the exterior, or *F* is on the exterior of T_j and all other edges from *j* are in the interior of T_j . In either case, $y_j(C, A) = q + 1$ for all *j* on c.

For the converse, suppose $y_j(C, A) = y_k(C, A) = q + 1$ for all *j*, *k* on *c* and some *q* on *A*. Then for each *j* on *c*, the triangle $T_j = (j, q, q + 1)$ is the smallest triangle

10



FIGURE 5 Depiction of special y-inversion [Color figure can be viewed at wileyonlinelibrary.com]

(by containment) from *j* to A that encloses B. By the minimality of T_j , no edge from *j* in D^* can cross the interior of T_j . The intersection of these triangles over all *j* on c is a face *F* of D^* containing B and adjacent to A at $\{q, q + 1\}$. By the observation, $x_i(A, C) = y_i(A, C)$ for all *i* on A.

For vertices *x* and *y* on a circle with *n* vertices, we define

$$\min d_n(x, y) \coloneqq \min\{d_n(x, y), d_n(y, x)\}.$$

The following lemma analyzes the situation when a term counting bichromatic crossings is minimized.

Lemma 11. Consider a tripartite-circle drawing with circles *A*, *B*, and *c* with *c* vertices on *c*. If $f_c(y_i(A, C), y_j(B, C)) = \min_{(u,v) \in [c]^2} f_c(u, v)$ for every $i \in A$ and $j \in B$, then there are vertices $u_A, u_B \in C$ with the following properties:

- (i) For both circles $D \in \{A, B\}$, $y_i(D, C) \in \{u_D, u_D + 1\}$ for all $i \in D$.
- (ii) For some circle $D \in \{A, B\}$, $y_i(D, C) = u_D$ for all $i \in D$. If c is even, then for both circles $D \in \{A, B\}$, $y_i(D, C) = u_D$ for all $i \in D$.

Proof. Suppose that $f_c(y_i(A, C), y_j(B, C))$ equals the minimum value of f_c for every $i \in A$ and $j \in B$. By Lemma 8, it holds that $|y_i(A, C) - y_j(B, C)| \in \{\lfloor c/2 \rfloor, \lceil c/2 \rceil\}$ and min $d_c(y_i(A, C), y_i(B, C)) = \lfloor c/2 \rfloor$.

First we prove property (i). Without loss of generality, we assume that D = A. Let $v := y_j(B, C)$ for some $j \in B$. The vertex v on c can have only one vertex at distance $\lfloor c/2 \rfloor = c/2$ if c is even and only two vertices at distance $\lfloor c/2 \rfloor$, adjacent to one another, if

11

Wh fy

c is odd. Since min $d_c(y_i(A, C), v) = \lfloor c/2 \rfloor$ for all $i \in A$, we get that the $y_i(A, C)$ labels are all the same in the even case, or all on one of two adjacent vertices in the odd case.

Next we prove property (ii). Suppose for contradiction that $y_i(A, C) = u_A, y_j(A, C) = u_A + 1, y_k(B, C) = u_B$, and $y_\ell(B, C) = u_B + 1$ for some $i, j \in A$ and $k, \ell \in B$. Since min $d_c(y_i(A, C), y_j(B, C)) = \lfloor c/2 \rfloor$ for every $i \in A$ and $j \in B$, we get that min $d_c(u_A, u_B) = \min d_c(u_A, u_B + 1) = \lfloor c/2 \rfloor$. No vertex on c other than u_A is at distance $\lfloor c/2 \rfloor$ from both u_B and $u_B + 1$, but $u_A + 1 \neq u_A$ must also be at distance $\lfloor c/2 \rfloor$ from both.

Now, we use these insights to improve the lower bound of Theorem 1.

Corollary 12 (Improvement of lower bound of Theorem 1). Let $\mathfrak{t} := \{(m, n, p), (n, p, m), (p, m, n)\}$. Then, for any integers $m, n, p \ge 3$, it holds that

$$\operatorname{cr}_{\odot}(K_{m,n,p}) \geq \sum_{(a,b,c)\in\mathfrak{t}} \left(\operatorname{cr}_{\odot}(K_{a,b}) + ab \left\lfloor \frac{c}{2} \right\rfloor \left\lfloor \frac{c-1}{2} \right\rfloor \right) + 2.$$

Proof. Recall that the lower bound in Theorem 1 was obtained by simultaneously minimizing all six terms in the formula of Theorem 7. Suppose a bichromatic crossing count $\sum_{j\in B} f_c(y_i(A, C), y_j(B, C))$ attains its minimum. Lemma 11(ii) implies without loss of generality that all $y_i(A, C)$ labels are equal. Lemma 10 then implies that $x_j(C, A) = y_j(C, A)$ for all j on c (and $x_j(C, B) = y_j(C, B)$ for all $j \in C$ if c is even). To achieve the minimum number of monochromatic crossings between A and c, the $x_j(C, A)$ labels must be equally spaced around A as already observed in [27]. Then, since $y_j(C, A) = x_j(C, A)$, the $y_j(C, A)$ labels are also equally spaced. Since $a \ge 3$, the $y_j(C, A)$ labels are on more than two points. By Lemma 11, the term $\sum_{\substack{i\in B\\j\in C}} f_a(y_i(B, A), y_j(C, A))$ does not attain its minimum.

For *c* even, Lemma 11(ii) further implies that all the $y_i(B, C)$ labels are equal. By Lemma 10, $x_j(C, B) = y_j(C, B)$ for all $j \in C$. If the minimum number of monochromatic crossings between B and c is achieved, then the bichromatic crossings term $\sum_{\substack{i \in A \\ j \in C}} f_b(y_i(A, B), y_j(C, B))$ also does not attain its minimum. By Lemma 8, if *c* is odd then at least one of the six terms is at least 2 more than its minimum. If *c* is even then at least two of the six terms are at least 1 more than their minima. Regardless of the parity of *c*, the lower bound given by minimizing all six terms simultaneously can be improved by 2.

3.2 Upper bounds

In this subsection, we provide drawings that settle the upper bounds of Theorem 1 and Corollary 2. We define the drawing within a small stripe around the equator of the sphere and visualize it by a rectangle where the left and right boundaries are identified. Consider Figure 6 for an illustration. In contrast to before, in the following drawings the interiors of the three circles are disjoint.



13

FIGURE 6 Illustration of the construction of a tripartite-circle drawing of $K_{m,n,p}$ with m = 4, n = 5, p = 6on a small strip around the equator of a sphere, the left and right boundaries of the rectangle are identified. The subdrawing induced by the circles A := M and B := N is highlighted by colors [Color figure can be viewed at wileyonlinelibrary.com

Definition (Linear Description). We start by defining the subdrawing D' induced by the vertices of two distinct circles A and B. Let A be a circle with a vertices and B a disjoint circle with b vertices. We assume that A is strictly left of B. The vertices are placed on the circles such that $\lceil a/2 \rceil$ and $\lceil b/2 \rceil$ vertices lie in the closed top halves of A and B, respectively; the vertices on A are labeled clockwise by $\{1, ..., a\}$ starting with the clockwise first vertex in the closed top half, while the vertices on B are labeled counterclockwise by $\{1, ..., b\}$ starting with the counterclockwise first vertex in the closed top half. Let ℓ_1 and ℓ_2 be two vertical lines separating A and B where ℓ_1 is strictly left of ℓ_2 .

- (i) On ℓ_1 we mark $a \cdot b$ points, which are labeled by $a_{i,j}$ for $i \in [a]$ and $j \in [b]$ such that the indices increase lexicographically from top to bottom. Each $a_{i,j}$ belongs to an edge of vertex i on A; between vertex i and $a_{i,j}$ the edge is realized by some x- and y-monotone curve $e_{i,i}^1$. Moreover, no two curves $e_{i,i}^1$ intersect.
- (ii) On ℓ_2 we mark $a \cdot b$ points, which are labeled by $b_{i,i}$ for $i \in [b]$ and $j \in [a]$ such that the indices increase lexicographically from top to bottom. Each $b_{i,j}$ belongs to an edge of vertex i on B; the edge between these two points is realized by some x- and y-monotone curves $e_{i,j}^2$. Moreover, no two curves $e_{i,j}^2$ intersect.
- (iii) Between ℓ_1 and ℓ_2 , we connect $a_{i,j}$ and $b_{j,i}$ by a straight-line segment.

The drawing D is obtained by constructing a drawing D' for each pair of circles and overlaying them. By construction, the drawing D has the following properties:

- (i) each edge is x-monotone,
- (ii) the drawing is partitioned into six vertical stripes; within each stripe every edge is x- and y-monotone,
- (iii) there exist two types of stripes, either containing AB/AB-crossings or AB/BC-crossings, and
- (iv) each edge is contained in three stripes.

These properties imply the following fact.

Proposition 13. The drawing D is simple.

Proof. For each pair of edges, there exists a unique stripe where the two edges potentially cross. Since, by property (ii), the edges are x- and y-monotone within each stripe, any pair of edges have at most one point in common. Because edges are x-monotone, no edge crosses itself. Thus, D is a simple drawing.

It remains to analyze the number of crossings for which we present two proofs. Proof 1 uses the fact that the drawings are simple together with Lemma 5 and 6 and offers an insight on what is needed to improve the construction. The crossing number can also be directly computed, as shown in Proof 2, providing an upper bound even if the drawings were not simple.

Proposition 14. For any integers $m, n, p \ge 3$, let $\mathfrak{t} := \{(m, n, p), (n, p, m), (p, m, n)\}$. The number of crossings in the drawing D is

$$\sum_{(a,b,c)\in\mathfrak{t}} \left(\binom{a}{2} \binom{b}{2} + ac \left\lfloor \frac{b}{2} \right\rfloor \left\lfloor \frac{b-1}{2} \right\rfloor \right).$$

Proof 1. It is easy to see from the construction that $x_i(A,B) = y_i(A, B), x_i(B, A) = y_i(B, A)$, and $d_b(y_i(A, B), y_j(C, B)) = \lfloor \frac{b}{2} \rfloor$. Consequently, by Lemma 5, the number of crossings of type AB/AB is

$$\sum_{1 \le i < j \le a} f_b(x_i(\mathbf{A}, \mathbf{B}), x_j(\mathbf{A}, \mathbf{B})) = \binom{a}{2} \binom{b}{2}.$$

By Lemma 6, the number of types AB/BC is

$$\sum_{\substack{1 \le i \le a \\ 1 \le j \le c}} f_b(y_i(\mathbf{A}, \mathbf{B}), y_j(\mathbf{C}, \mathbf{B})) = ac \left(\left(\frac{\lfloor b/2 \rfloor}{2} \right) + \left(\frac{\lceil b/2 \rceil}{2} \right) \right) = ac \lfloor b/2 \rfloor \lfloor (b-1)/2 \rfloor.$$

This finishes the first proof.

Proof 2. Alternatively, we count the number of crossings directly. By definition, the AB/AB crossings occur between ℓ_1 and ℓ_2 ; in this part of the drawing the edges are straight-line segments. Any pair of vertices on circle A and any pair of vertices on circle B together form exactly one crossing. We have $\binom{a}{2}\binom{b}{2}$ crossings.

For the crossings of type AB/BC, it suffices to count the bundle crossings. If two bundles cross they add *ac* crossings. Moreover, it follows from the construction that two bundles cross if and only if they are both in the top or both in the bottom half. Consequently, the number of crossings is

⊥WILEY-

$$ac\left(\left(\frac{\lfloor b/2 \rfloor}{2}\right) + \left(\frac{\lceil b/2 \rceil}{2}\right)\right).$$

As shown above, this evaluates to $ac\lfloor b/2 \rfloor \lfloor (b-1)/2 \rfloor$ and therefore finishes the second proof of the proposition.

As the proposition was the last missing item, this finishes the proof of the upper bound and thus of Theorem 1. Note that this construction achieves the minimum possible number of bichromatic crossings by Lemma 9.

3.3 | Balanced case

Theorem 1 and Corollary 12 imply Corollary 2 for the special case of m = n = p.

Proof of Corollary 2. For the lower bound, Theorem 1 and Corollary 12 give that

$$\operatorname{cr}_{\odot}(K_{n,n,n}) \ge 3\operatorname{cr}_{\odot}(K_{n,n}) + 3n^2 \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor + 2.$$

With the bipartite cylindrical crossing number from Equation (1) we have

$$\operatorname{cr}_{\mathfrak{I}}(K_{n,n,n}) \geq 3n\binom{n}{3} + 3n^2 \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor + 2.$$

For the upper bound, the construction includes drawings for $K_{n,n,n}$. In this case, we obtain highly symmetric drawings, which are especially appealing. In particular, such a drawing can be defined by two consecutive stripes; see Figure 9C. The formula simplifies to

$$3\binom{n}{2}^2 + 3n^2 \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-1}{2} \right\rfloor.$$

While the lower bound order is ${}^{5}/_{4} \cdot n^{4}$, the upper bound order is ${}^{6}/_{4} \cdot n^{4}$. Consequently, the bounds are fairly close. Moreover, instead of a linear representation, similar drawings can be defined in a *cyclic* way, as shown in Figure 7.

Remark. By a slight modification, we improve the upper bound. To do so, we place at least one vertex on the intersection of the closed top and bottom half of the circle and route half of its incident edges via the upper half and the other half of its edges via the bottom half. This idea is used to construct the drawings in Figure 8 (and Figure 9C).

Depending on the parity of *n*, the number of monochromatic crossings between two circles is

WILFY



16





FIGURE 8 Improved tripartite-circle drawings of $K_{n,n,n}$ for n = 5,6 [Color figure can be viewed at wileyonlinelibrary.com]

$$\begin{cases} \binom{n-1}{2}\binom{n}{2} + (n-1)\left(\binom{\lfloor n/2 \rfloor}{2} + \binom{\lceil n/2 \rceil}{2}\right) & n \text{ odd,} \\ 2\left(\binom{n/2-1}{2}n^2 + \frac{1}{2}(n-2)n^2 + \frac{1}{4} \cdot n^2\right) & n \text{ even,} \end{cases}$$

while the number of bichromatic crossings is

$$\begin{cases} n^2 \left(\binom{\lfloor n/2 \rfloor}{2} + \binom{\lceil n/2 \rceil}{2} \right) & n \text{ odd,} \\ \binom{(n-1)}{2}^2 + 2(n-1) \left(\binom{n/2-1}{2} + \binom{n/2}{2} \right) + (n/2)^2 + (n/2-1)^2 & n \text{ even.} \end{cases}$$

Consequently, multiplying by three and summing both terms, the number of crossings evaluates to

$$\begin{cases} \frac{3}{4}(2n^4 - 5n^3 + 3n^2 + n - 1) & n \text{ odd,} \\ \frac{3}{4}(2n^4 - 6n^3 + 7n^2) & n \text{ even.} \end{cases}$$

Unfortunately, this improves only lower-order terms, that is, the number of saved crossings is



FIGURE 9 Drawings of $K_{n,n,n}$ for n = 2, 3, 4 with few crossings. (B, C) The two drawings are obtained by considering either the dash-dotted edge or the dotted edge [Color figure can be viewed at wileyonlinelibrary.com]

	Lower bound	Improved	Improved	Upper bound
n	Corollary 2	lower bound	upper bound	Corollary 2
2	-	3	3	-
3	38	-	42	54
4	146	147	175	204
5	452	-	528	600
6	1010	-	1161	1323
7	2060	-	2430	2646
8	3650	-	4176	4656
9	6158	-	7296	7776
10	9602	-	11,025	12,075



FIGURE 10 A 3-circle drawing of $K_{4,4,4}$ with 175 crossings [Color figure can be viewed at wileyonlinelibrary.com]

$$\begin{cases} \frac{3}{4}(n^3 - n^2 - n + 1) & n \text{ odd,} \\ \frac{3}{2}(n^3 - 3n^2) & n \text{ even.} \end{cases}$$

3.3.1 | Balanced case with few vertices

 \perp WILEY

In this section, we present numerical results and improved drawings of $K_{n,n,n}$ for small values of *n*. The values are summarized in Table 1. We improve the upper bounds with concrete drawings and the lower bounds with Corollary 12 and the following fact:

Proposition 15. For any integers m, n, and p,

$$\operatorname{cr}_{\odot}(K_{m,n,p}) \ge cr(K_{m+n+p}) - \binom{m}{4} - \binom{n}{4} - \binom{p}{4}.$$

The proposition follows from the fact that a tripartite-circle drawing of $K_{m,n,p}$ yields a drawing of the complete graph K_{m+n+p} by adding all straight-line segments within the three circles; see also Section 4.

In the following we explain how to obtain the bounds displayed in Table 1. For n = 2, it holds that $\operatorname{cr}_{\odot}(K_{2,2,2}) \ge 3$ since $cr(K_6) = 3$ and Figure 9A shows that three crossings can be attained.

For n = 3, a lower bound of 38 follows from Corollary 12 and the upper bound of 42 from the drawing in Figure 9B.

In case n = 4, we use Proposition 15 for the lower bound. Since $cr(K_{12}) = 150$, we obtain $cr_{\odot}(K_{4,4,4}) \ge cr(K_{12}) - 3 = 147$. For the upper bound, Figure 10 presents a drawing with 175 crossings.

This drawing is obtained by a slight modification of the drawing corresponding to Figure 9C with the dash–dotted edge. In particular, in the middle copy (orange), the long edge between the leftmost vertex of $_{\rm B}$ and the rightmost vertex of $_{\rm C}$ is drawn in the top half, while its corresponding edges in the other two copies are drawn in the bottom half. This saves the two crossings between the middle long edge and the left and right long edges.

For $n \ge 5$, we use the ideas of Section 3.3 to improve the upper bounds.

4 | CONNECTION TO THE HARARY-HILL CONJECTURE

The Harary–Hill Conjecture [11,20] states that the number of crossings in any drawing (in the plane) of the complete graph K_n is at least

$$H(n) \coloneqq \frac{1}{4} \left[\frac{n}{2} \right] \left[\frac{n-1}{2} \right] \left[\frac{n-2}{2} \right] \left[\frac{n-3}{2} \right].$$

Drawings with exactly H(n) crossings [12,20] show that $cr(K_n) \le H(n)$. The Harary–Hill conjecture has been confirmed for $n \le 12$ (see [19] for $n \le 10$ and [25] for n = 11, 12), and either $cr(K_{13}) = H(13)$ or H(13) - 2 [1]. The conjecture has been proved when restricted to certain families of graphs [2–5,10,24].

For decades, only two families of drawings of K_n with H(n) crossings were known, shown in Figure 11: the Blažek–Koman construction [12], which is an instance of a 1-circle drawing, and the Harary–Hill construction [20], which is an instance of a balanced restricted 2-circle drawing.

Ábrego et al. announced in [6] a new family of drawings of K_n having the property that each edge is crossed at least once, so these are not k-circle drawings. Kynčl and others [22] naturally asked about the existence of alternative k-circle constructions of K_n with H(n)crossings: Is $H(n) = bcr_{\otimes}(K_n)$ for some $k \ge 3$? (Recall that bcr_{\otimes} denotes the minimum number of crossings in a balanced restricted k-circle crossing drawing.) Figure 12 shows



Construction of Blazek and Koman a 1-circle drawing.

Construction of Harary and Hill; a 2-circle drawing.





FIGURE 12 Balanced 1-, 2-, and 3-circle drawings of K_6 , each with 3 = H(6) crossings [Color figure can be viewed at wileyonlinelibrary.com]

crossing-optimal balanced 1-, 2-, and 3-circle drawings of K_6 . We prove that balanced restricted 3-circle drawings are suboptimal for n large enough.

Corollary 16. For n = 9, 10 and $n \ge 13$, the number of crossings in any balanced restricted 3-circle drawing of K_n exceeds H(n), that is, $bcr_{\odot}(K_n) > H(n)$.

Proof. Suppose $n \ge 14$. Let $q \ge 5$ and $r \in \{-1, 0, 1\}$ be the unique integers such that n = 3q + r. We want to show that $bcr_{\odot}(K_n) - H(n) > 0$. Consider a balanced restricted 3-circle drawing of K_n with q, q, and q + r vertices on the three circles. Then

$$\operatorname{bcr}_{\odot}(K_n) = \operatorname{cr}_{\odot}(K_{q,q,q+r}) + 2\binom{q}{4} + \binom{q+r}{4}.$$
(4)

We use Theorem 1 to bound $\operatorname{cr}_{\odot}(K_{q,q,q+r})$ and $\lfloor q/2 \rfloor \lfloor (q-1)/2 \rfloor \ge q(q-2)/4$ to remove the floor function, which yields

$$\operatorname{cr}_{\odot}(K_{q,q,q+r}) \ge \operatorname{cr}_{\odot}(K_{q,q}) + 2\operatorname{cr}_{\odot}(K_{q,q+r}) + q^{2} \left\lfloor \frac{q+r}{2} \right\rfloor \left\lfloor \frac{q+r-1}{2} \right\rfloor + 2q(q+r) \left\lfloor \frac{q}{2} \right\rfloor \left\lfloor \frac{q-1}{2} \right\rfloor$$
$$\ge \operatorname{cr}_{\odot}(K_{q,q}) + 2\operatorname{cr}_{\odot}(K_{q,q+r}) + \frac{1}{4}q^{2}(q+r)(3q+r-6).$$

By Equation (1), we have $\operatorname{cr}_{\otimes}(K_{q,q}) = q\binom{q}{3}$. By Equation (2), we obtain $\operatorname{cr}_{\otimes}(K_{q,q-1}) = (q-2)\binom{q}{3}$ and $\operatorname{cr}_{\otimes}(K_{q,q+1}) = (q-1)\binom{q+1}{3}$. Thus

$$\operatorname{cr}_{\textcircled{0}}(K_{q,q,q+r}) \geq \begin{cases} 3q\binom{q}{3} + \frac{1}{4}q^{3}(3q-6) & \text{if } r = 0, \\ (3q-4)\binom{q}{3} + \frac{1}{4}q^{2}(q-1)(3q-7) & \text{if } r = -1, \\ q\binom{q}{3} + 2(q-1)\binom{q+1}{3} + \frac{1}{4}q^{2}(q+1)(3q-5) & \text{if } r = 1. \end{cases}$$
(5)

The result holds for $n \ge 14$ $(q \ge 5)$ by (4), (5), and $H(n) \le \frac{1}{64}(n-1)^2(n-3)^2$:

$$\operatorname{bcr}_{\odot}(K_n) - H(n) \ge \begin{cases} \frac{1}{64}(7q^4 - 24q^3 - 46q^2 + 24q - 9) > 0 & \text{if } r = 0, \\ \frac{q}{192}(21q^3 - 100q^2 - 36q + 112) > 0 & \text{if } r = -1, \\ \frac{q}{192}(q + 2)(21q^2 - 86q - 8) > 0 & \text{if } r = 1. \end{cases}$$

Finally, by (4) and Corollary 12, $bcr_{\odot}(K_9) = cr_{\odot}(K_{3,3,3}) \ge 38 > 36 = H(9)$, $bcr_{\odot}(K_{10}) = cr_{\odot}(K_{3,3,4}) + 1 \ge 64 > 60 = H(10)$, and $bcr_{\odot}(K_{13}) = cr_{\odot}(K_{4,4,5}) + 7 \ge 229 > 225 = H(13)$.

Our previous argument does not settle the cases n = 8, n = 11 and 12. The Harary–Hill constructions for $n \le 5$ are in fact balanced restricted 3-circle drawings, and we give balanced restricted 3-circle drawings of K_6 and K_7 in Figures 12C and 13, respectively. If we allow for unbalanced constructions for $n \ge 8$, the crossing number of $\operatorname{cr}_{\odot}(K_{2,2,n-4})$ given in Equation (3) implies that

20



FIGURE 13 Balanced 3-circle drawing of K_7 with 9 = H(7) crossings [Color figure can be viewed at wileyonlinelibrary.com]

$$\operatorname{cr}_{O3}(K_{2,2,n-4}) + \binom{n-4}{4} = \binom{n-4}{4} + \frac{3}{2}(n-4)^2 - (n-4) - \begin{cases} 3 & n \text{ even} \\ 3/2 & n \text{ odd} \end{cases} \ge H(n),$$

with equality if and only if $8 \le n \le 11$. That is, for $n \ge 8$ the drawings of K_n given by a crossingoptimal 3-circle drawing of $K_{2,2,n-4}$ together with the straight-line drawings inside of the three circles achieve H(n) crossings if and only if $8 \le n \le 11$.

5 | CONCLUSION AND OPEN PROBLEMS

In this paper, we prove upper and lower bounds on the tripartite-circle crossing number of complete tripartite graphs. For the lower bound, we introduce formulas describing the number of crossings in a tripartite-circle drawing. For the upper bounds, we present drawings. While there exist restricted balanced 2-circle drawings achieving the Harary–Hill bound, our results imply that this is not the case for balanced restricted 3-circle drawings for $n \ge 13$. It remains open for future work whether the same holds for *k*-partite circle drawings when k > 3. We have made progress in the direction of extending our work to k > 3 and plan to return to this question in a subsequent paper. We conclude with a list of interesting open problems for future work:

- Do there exist *k*-circle drawings achieving the Harary-Hill bound for k > 3?
- Can the number of crossings of *k*-circle drawings generally be described by labels analogous to *x*-and*y*-labels?
- How are crossing-minimal k-circle drawings characterized?
- What are the exact values for small graphs? Is $cr_{\odot}(K_{3,3,3}) = 42$? For the remaining displayed values in Table 1, we believe that the truth lies closer to the presented upper bounds. In particular, it remains to develop better tools to improve the lower bounds.
- Are there balanced restricted 3-circle drawings achieving the Harary–Hill bound for K_8 , K_{11} , or K_{12} ?
- We know that balanced restricted 3-circle drawings do not (in general) achieve the Harary–Hill bound, and that extending tripartite-circle drawings of $K_{2,2,n-4}$ does not (in general) achieve the Harary–Hill bound. Are there other unbalanced restricted 3-circle drawings that do achieve the Harary–Hill bound?

ACKNOWLEDGMENTS

As this project started at the MRC workshop "Beyond Planarity: Crossing Numbers of Graphs," we thank the organizers and all participants for the fruitful atmosphere. We thank

WILFY

CAMACHO ET AL.

Julia Böttcher for helpful feedback on a draft of this manuscript. Open Access funding enabled and organized by Projekt DEAL.

ORCID

Linda Kleist D https://orcid.org/0000-0002-3786-916X

REFERENCES

- B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, T. Hackl, J. Pammer, A. Pilz, P. Ramos, G. Salazar, and B. Vogtenhuber, *All good drawings of small complete graphs*, Proc. 31st European Workshop on Computational Geometry (EuroCG), 2015, pp. 57–60.
- B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, D. McQuillan, B. Mohar, P. Mutzel, P. Ramos, R. B. Richter, and B. Vogtenhuber, *Bishellable drawings of K_n*, SIAM J. Discrete Math. **32** (2018), no. 4, 2482–2492.
- 3. B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, P. Ramos, and G. Salazar, *The 2-page crossing number of K_n*, Discrete Comput. Geom. **49** (2013), no. 4, 747–777.
- B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, P. Ramos, and G. Salazar, More on the crossing number of K_n: Monotone drawings, Electron. Notes Discrete Math. 44 (2013), 411–414.
- 5. B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, P. Ramos, and G. Salazar, *Shellable drawings and the cylindrical crossing number of K_n*, Discrete Comput. Geom. **52** (2014), no. 4, 743–753.
- 6. B. M. Ábrego, O. Aichholzer, S. Fernández-Merchant, P. Ramos, and B. Vogtenhuber, *Non-shellable drawings of K_n with few crossings*, Proc. 26th Annual Canadian Conference on Comp. Geom. (CCCG), 2014.
- 7. B. M. Ábrego, S. Fernández-Merchant, and A. Sparks, *The cylindrical crossing number of the complete bipartite graph*, Graphs Combin. **36** (2019), 205–220.
- 8. K. Asano, *The crossing number of* K_{1,3,n} and K_{2,3,n}, J. Graph Theory. **10** (1986), no. 1, 1–8.
- 9. C. Bachmaier, A radial adaptation of the Sugiyama framework for visualizing hierarchical information, IEEE Trans. Vis. Comput. Graph. **13** (2007), no. 3, 583–594.
- 10. M. Balko, R. Fulek, and J. Kynčl, Crossing numbers and combinatorial characterization of monotone drawings of K_n , Discrete Comput. Geom. 53 (2015), no. 1, 107–143.
- 11. L. Beineke and R. Wilson, *The early history of the brick factory problem*, Math. Intelligencer. **32** (2010), no. 2, 41–48.
- 12. J. Blažek, and M. Koman, *A minimal problem concerning complete plane graphs*, Theory of Graphs and its Applications, M. Fiedler (ed.), Czech. Acad. of Sci., Prague, 1964, pp. 113–117.
- C. Camacho, S. Fernández-Merchant, R. Kirsch, L. Kleist, E. Matson, M. J. Milutinović, and J. White, *The tripartite-circle crossing number of K*_{2,2,n}, arxiv preprint: 2108.01032, 2021.
- F. Duque, H. González-Aguilar, C. Hernández-Vélez, J. Leaños, and C. Medina, *The complexity of computing the cylindrical and the t-circle crossing number of a graph*, Electron. J. Combin. 25 (2018), no. 2, P2–43.
- 15. M. R. Garey and D. S. Johnson, *Crossing number is NP-complete*, SIAM J. Algebr. Discrete Methods. 4 (1983), no. 3, 312–316.
- 16. E. Gethner, L. Hogben, B. Lidickỳ, F. Pfender, A. Ruiz, and M. Young, On crossing numbers of complete tripartite and balanced complete multipartite graphs, J. Graph Theory. 84 (2017), no. 4, 552–565.
- 17. M. Ginn and F. Miller, The crossing number of $K_{3,3,n}$, Congr. Numer. 221 (2014), 49–54.
- 18. R. K. Guy, A combinatorial problem, Nabla (Bull. Malayan Math. Soc.). 7 (1960), 68–72.
- 19. R. K. Guy, *Crossing numbers of graphs*, Graph Theory and Applications, Springer, Y. Alavi, D. R. Lick, and A. T. White (eds.), Berlin, 1972, pp. 111–124.
- F. Harary and A. Hill, On the number of crossings in a complete graph, Proc. Edinb. Math. Soc. 13 (1963), no. 4, 333–338.
- 21. P. Hliněngrý, Crossing number is hard for cubic graphs, J. Combin. Theory Ser. B. 96 (2006), no. 4, 455–471.
- 22. J. Kynčl, Drawings of complete graphs with Z(n) crossings. https://mathoverflow.net/questions/128878
- 23. F. T. Leighton, Complexity issues in VLSI, MIT Press, Cambridge, MA, United States, 1983.
- 24. P. Mutzel and L. Oettershagen, *The crossing number of semi-pair-shellable drawings of complete graphs*, Proc. 30th Annual Canadian Conference on Comp. Geom. (CCCG) 2018, pp. 11–17.

22

- 25. S. Pan and R. B. Richter, *The crossing number of K*₁₁ *is 100*, J. Graph Theory. **56** (2007), no. 2, 128–134.
- 26. M. J. Pelsmajer, M. Schaefer, and D. Štefankovic, *Odd crossing number and crossing number are not the same*, Discrete Comput. Geom. **39** (2008), 442–454.
- 27. R. B. Richter and C. Thomassen, *Relations between crossing numbers of complete and complete bipartite graphs*, Amer. Math. Monthly. **104** (1997), no. 2, 131–137.
- 28. M. Schaefer, *The graph crossing number and its variants: A survey*, Electron. J. Combin. **DS21** (2021). https://doi.org/10.37236/2713
- 29. K. Zarankiewicz, On a problem of P. Turán concerning graphs, Fund. Math. 1 (1955), no. 41, 137-145.

How to cite this article: C. Camacho, S. Fernández-Merchant, M. Jelić Milutinović, R. Kirsch, L. Kleist, E. B. Matson, and J. White, *Bounding the tripartite-circle crossing number of complete tripartite graphs*, J. Graph Theory. (2021), 1–23. https://doi.org/10.1002/jgt.22763